

AN INDEX THEOREM FOR GENERAL EQUILIBRIUM MODELS WITH PRODUCTION¹

BY TIMOTHY J. KEHOE²

In this paper we prove a global index theorem for general equilibrium models with activity analysis production technologies. We begin by constructing a single-valued function whose fixed points are equivalent to the equilibria of such a model. We then associate each fixed point with an index that is an integer determined by the local properties of this function at that point. The global index theorem makes a statement about the sum of all the indices of equilibria that implies conditions sufficient for uniqueness of equilibrium.

1. INTRODUCTION

IT IS WELL KNOWN that some variant of Brouwer's fixed point theorem can be used to prove the existence of equilibrium prices for a general model of economic competition. However, simple existence proofs leave many questions unanswered: For example, is the equilibrium price vector unique? If not, is it locally unique? Does it vary continuously with the underlying parameters of the model? Answers to such questions are intimately linked with this model's applicability to problems of comparative statics. Consequently, it is not surprising to find that a large amount of effort has been devoted to analyzing these questions. To provide complete answers, however, requires a more detailed set of assumptions than those required for simple existence proofs. For this reason we adopt a differentiable framework.

There have been many approaches to answering the question of when an equilibrium is unique (see Arrow and Hahn [1, Chapter 9] for a survey). Several recent approaches have made use of a tool borrowed from the field of algebraic topology, the fixed point index (Dierker [6, 7, 8], Mas-Colell [23, 24], and Varian [32]). Researchers have recognized the close connection between this tool, when cast in a differentiable framework, and the concept of regularity introduced by Debreu [2] in response to the questions of local uniqueness and continuity. With the notable exceptions of Fuchs [13], Mas-Colell [22, 23, 24], and Smale [27], researchers have focused their attention on the pure exchange model that allows no production. One reason for this focus has been that, in contrast to the pure exchange model, where the excess demand function has been the natural subject of study, the model with production has had no obvious, single-valued, differentiable function to investigate.

¹ The research presented here was partially funded through NSF Grant No. SOC77-00210. Earlier versions of this paper were presented at the NBER Conference-Seminar on the Theory of General Economic Equilibrium, University of California, Berkeley, May, 1979, and at the North American Meeting of the Econometric Society, University of Montreal, June, 1979.

² I am grateful to my dissertation advisors Herbert Scarf and Andreu Mas-Colell for their constant advice, encouragement, and guidance. It was Professor Scarf who first suggested the idea for this paper. Both he and Professor Mas-Colell have helped me see it through numerous corrections and revisions.

Using a variant of a device due to Todd [31], we define a single-valued function whose fixed points are equivalent to equilibria of a model with an activity analysis production technology. We associate each fixed point of this function with an index that is an integer determined by the local properties of the function at that point. The global index theorem makes a statement about the sum of all the indices of equilibria allowing us to establish conditions sufficient for uniqueness.

For "almost all" economies with differentiable consumer demand functions the index of an equilibrium price vector is either +1 or -1 depending on the sign of

$$\det \begin{bmatrix} -\bar{J} & -\bar{B} \\ \bar{B}' & 0 \end{bmatrix}.$$

Here \bar{J} is formed by deleting the first row and column from the Jacobian matrix of demand functions; \bar{B} is formed by deleting the first row from the matrix of activities in use at equilibrium. The global index theorem states that the sum of all the indices of equilibrium of an economy is equal to +1. Mas-Colell has independently derived many of the results presented in this paper without directly utilizing such a single-valued function. His approach, however, involves mathematical machinery more complex than that employed here.

2. THE MODEL

Let us begin by describing a simple version of the Walrasian model of economic equilibrium. We assume that there is a finite number, n , of perfectly divisible commodities. On the consumption side of the model responses of consumers to a vector of nonnegative prices $\pi = (\pi_1, \dots, \pi_n)$ are aggregated into a vector of market excess demand functions $\xi(\pi) = (\xi_1(\pi), \dots, \xi_n(\pi))$. We take these functions to be completely arbitrary except for the following assumptions:

ASSUMPTION 1 (Differentiability): Each ξ_i is a continuously differentiable function defined, for the sake of simplicity, over the domain of all nonnegative prices except the origin, $\mathbb{R}_+^n \setminus \{0\}$.

ASSUMPTION 2 (Homogeneity): Each ξ_i is homogeneous of degree zero; that is, $\xi_i(t\pi) = \xi_i(\pi)$ for all $t > 0$.

ASSUMPTION 3 (Walras's law): The vector function ξ obeys Walras's law, $\pi' \xi(\pi) = 0$.

Several comments about these assumptions are in order. The homogeneity assumption (Assumption 2) embodies the understanding that only relative prices matter in a model where money plays no explicit role. Walras's law (Assumption 3) dictates that all income is spent. These two assumptions are standard and completely general; the differentiability assumption (Assumption 1) is more specific. It is this assumption, however, that is central to our differentiable

approach to the study of equilibrium. Recent studies have suggested that assuming excess demand to be a continuously differentiable, single-valued function, rather than the more general upper-semi-continuous, set-valued correspondence, is not overly restrictive (Debreu [3, 4] and Mas-Colell [21]). The central point of these studies is that arbitrarily small perturbations in the underlying preferences of individual consumers suffice to make aggregate excess demand a differentiable function. Since we use the excess demand function, rather than consumers' preferences, to specify the consumption side of the model, we shall not pursue this issue. We should note, however, that Assumption 1 is further restrictive in requiring ξ to be continuous even at points where some $\pi_i = 0$. The technique usually employed to ensure that this property holds is to put bounds on the consumption sets of individual consumers in a way that does not disturb the equilibria of the economy. Alternatively, it is a relatively straightforward matter to relax Assumption 1 utilizing only the concepts that we have defined here (see Kehoe [18], pp. 62-65). For the present, however, we shall ignore this minor technical point. The above assumptions completely describe the consumption side of the model. Indeed, if Assumption 1 is replaced by a weakened continuity assumption, then Assumptions 1-3 fully characterize aggregate excess demand functions under general conditions (McFadden, Mas-Colell, Mantel, and Richter [20]).

The production technology is specified by an activity analysis matrix A with n rows and m columns. Aggregate production is denoted by Ay where y is an $m \times 1$ vector of nonnegative activity levels. We assume that A satisfies the following assumptions:

ASSUMPTION 4 (Free disposal): A includes n free disposal activities, one for each commodity.

ASSUMPTION 5 (Boundedness): There are no outputs without any inputs, $\{Ay \in R^n | y \geq 0, Ay \geq 0\} = \{0\}$.

Alternative forms of these assumptions are useful: The free disposal assumption implies that the $n \times n$ matrix $-I$ is a submatrix of A . The boundedness assumption implies that there is some vector π , strictly positive, such that $\pi'A \leq 0$.

For our present purposes an economy is completely described by an excess demand function representing the consumption side and an activity analysis matrix representing the production side.

DEFINITION: An *equilibrium* of an economy (ξ, A) is a price vector $\hat{\pi}$ that satisfies the following conditions: (a) $\hat{\pi}'A \leq 0$. (b) There exists $\hat{y} \geq 0$ such that $\xi(\hat{\pi}) = A\hat{y}$. (c) $\sum_{i=1}^n \hat{\pi}_i = 1$.

The condition $\hat{\pi}'A \leq 0$ implies that at prices $\hat{\pi}$ no excess profits can be made. $\xi(\hat{\pi}) = A\hat{y}$ requires that demand equal supply at equilibrium. This condition, together with Walras's law, implies that $\hat{\pi}'A\hat{y} = 0$. Thus profits are maximized by

the aggregate production plan $A\hat{y}$. Since Assumption 1 rules the vector $\pi = 0$ out of consideration, we use the homogeneity assumption (Assumption 2) and the free disposal assumption (Assumption 4) to restrict our attention to the unit simplex

$$S = \left\{ \pi \in R^n \mid \pi_i \geq 0, \sum_{i=1}^n \pi_i = 1 \right\}$$

when examining equilibrium positions. At this stage it might seem more appropriate to include the vector of activity levels \hat{y} directly in the definition of equilibrium. However, we shall soon impose conditions on (ξ, A) that would make this addition superfluous.

We shall find it useful to consider whole spaces of economies. To do so, we must specify some topological structure on the space of economies satisfying Assumptions 1-3 and on the set of activity analysis matrices satisfying Assumptions 4 and 5. A topology on a space is specified by defining a system of open sets. Recall that in a metric space this is done by employing the concept of distance between two points in that space. We now give the space of economies the structure of a metric space: Let $\mathcal{A} \subset \{-I\} \times R^{n \times (m-n)}$ be the space of activity analysis matrices that satisfy Assumptions 4 and 5; here $-I$ is the submatrix of free disposal activities. We endow \mathcal{A} with the standard topology on $R^{n \times (m-n)}$ by defining the metric

$$d(A^1, A^2) = \left(\sum_{i=1}^n \sum_{j=n+1}^m (a_{ij}^1 - a_{ij}^2)^2 \right)^{\frac{1}{2}}$$

for any $A^1, A^2 \in \mathcal{A}$. Let \mathcal{D} be the space of C^1 functions satisfying Assumptions 1-3. We endow \mathcal{D} with the topology of uniform C^1 convergence on compacta. Letting M be some compact subset of $R^n \setminus \{0\}$, we define the metric

$$d_M(\xi^1, \xi^2) = \sup_{i, \pi \in M} |\xi_i^1(\pi) - \xi_i^2(\pi)| + \sup_{i, j, \pi \in M} \left| \frac{\partial \xi_i^1}{\partial \pi_j}(\pi) - \frac{\partial \xi_i^2}{\partial \pi_j}(\pi) \right|$$

for any $\xi^1, \xi^2 \in \mathcal{D}$. Since we adopt Assumptions 1 and 2, we shall use S as the compact domain on which maps ξ^1, ξ^2 are compared. (This topology is the same as the Whitney C^1 topology if we restrict the domain of our maps to the compact set S .) The space of economies $\mathcal{E} = \mathcal{D} \times \mathcal{A}$ has the induced product topology: For any $(\xi^1, A^1), (\xi^2, A^2) \in \mathcal{E}$ we define the metric $d[(\xi^1, A^1), (\xi^2, A^2)] = d_S(\xi^1, \xi^2) + d(A^1, A^2)$.

DEFINITION: An *economy* (ξ, A) is an element of the metric space $\mathcal{E} = \mathcal{D} \times \mathcal{A}$.

3. THE MAPPING

Let us now define a mapping of S into itself whose fixed points are equivalent to equilibria of an economy (ξ, A) . Letting N be any nonempty, closed, subset of R^n , we define the projection map $p^N: R^n \rightarrow N$ by the rule that associates any point $q \in R^n$ with the point $p^N(q)$ that is closest to q in terms of Euclidean distance. It is well known that p^N is continuous if N is convex. Observe that our definition of equilibrium implies that any equilibrium is an element of the nonempty, closed, convex set $S_A = \{\pi \in R^n \mid \pi'A \leq 0, \sum_{i=1}^n \pi_i = 1\} \subset S$.




FIGURE 1

DEFINITION: For any economy $(\xi, A) \in \mathcal{E}$, define the map $g : S \rightarrow S$ by the rule

$$g(\pi) = p^{S_A}(\pi + \xi(\pi)).$$




FIGURE 2

It will sometimes be convenient to refer to g as $g_{(\xi, A)}$.

Note that, since g is the composition of two continuous maps, it too is continuous. We shall apply Brouwer's fixed point theorem to this map g to prove the existence of an equilibrium price vector for g .

BROUWER'S FIXED POINT THEOREM: *If $f : M \rightarrow M$ is a continuous map of some nonempty, compact, convex set $M \subset \mathbb{R}^n$ into itself, then f leaves some point fixed; that is, there exists $\hat{x} \in M$ such that $\hat{x} = f(\hat{x})$.*

The motivation for our definition of g is clearly seen in the following theorem.

THEOREM 1: *Fixed points $\hat{\pi} = g(\hat{\pi})$ of the map g and equilibria of the economy $(\xi, A) \in \mathcal{E}$ are equivalent.*

PROOF: $\hat{\pi} = g(\hat{\pi})$ if and only if $\hat{\pi}$ is the unique solution to the quadratic programming problem

$$\begin{aligned} &\min \frac{1}{2}(p - \hat{\pi} - \xi(\hat{\pi}))'(p - \hat{\pi} - \xi(\hat{\pi})) \\ &\text{subject to } p'A \leq 0, \\ &\quad p'e = 1, \quad \text{where } e = (1, \dots, 1). \end{aligned}$$

By the Kuhn-Tucker theorem there exist nonnegative $\hat{y} = (\hat{y}_1, \dots, \hat{y}_m)$ and $\hat{\lambda}$ such that

$$p - \hat{\pi} - \xi(\hat{\pi}) + A\hat{y} + \hat{\lambda}e = 0$$

and $p'A\hat{y} = 0$. But $p = \hat{\pi}$ implies that $\xi(\hat{\pi}) = A\hat{y} + \hat{\lambda}e$. Applying Walras's law, we obtain

$$\begin{aligned} 0 &= \hat{\pi}'\xi(\hat{\pi}) \\ &= \hat{\pi}'A\hat{y} + \hat{\pi}'\hat{\lambda}e \\ &= \hat{\lambda}. \end{aligned}$$

Therefore, since $\hat{\pi} \in S_A$, the equilibrium conditions are satisfied. Q.E.D.

Consequently, Brouwer's theorem implies the existence of equilibrium and hence the logical consistency of our model. The restriction of the domain of g to S , while entirely natural for proving existence of equilibrium, would make our discussion of regularity and index theory awkward. We want to study the derivatives of g . To simplify matters, we define X to be a smooth (that is, C^1) n -dimensional manifold with boundary, embedded in \mathbb{R}^n so that it contains S in its interior and does not contain the origin. We now extend the domain of the map g to X .

LEMMA 1: *Let ξ be any function satisfying Assumption 1 and let $X \subset \mathbb{R}^n$ be defined as above. ξ can be extended to a C^1 map with domain X .*

PROOF: We demonstrate our contention by constructing a C^1 map $\xi^* : \mathbb{R}^n \rightarrow \mathbb{R}^n$ that satisfies $\xi^*(\pi) = \xi(\pi)$ for all $\pi \in X \cap \mathbb{R}_+^n$. There are two steps in this construction. First, we deal with the relatively minor technical problem of ξ being undefined at the origin. Second, we follow Saigal and Simon [26] in extending ξ to a C^1 map on all \mathbb{R}^n .

Since X is compact and does not contain $0 \in \mathbb{R}^n$, the continuous function $\|\pi\|$ achieves a minimum $\alpha > 0$ on X . Let β be such that $\|\pi\| \geq \alpha > \beta > 0$ for all $\pi \in X$. Take a smooth function $\theta : \mathbb{R}^n \rightarrow \mathbb{R}$ that satisfies

$$\begin{aligned} \theta(\pi) &= 1 && \text{if } \|\pi\| \leq \beta, \\ \theta(\pi) &= 0 && \text{if } \|\pi\| \geq \alpha, \\ 0 < \theta(\pi) < 1 && \text{if } \beta < \|\pi\| < \alpha. \end{aligned}$$

The construction of such a function, known as a bump function, is a standard exercise (see Hirsch [17, pp. 41-42]). We define $\xi_\beta^0(\pi) = (1 - \theta(\pi))\xi(\pi)$ for all $\pi \in \mathbb{R}_+^n \setminus \{0\}$ and $\xi_\beta^0(\pi) = 0$ for $\pi = 0$. Note that ξ_β^0 is C^1 , maps \mathbb{R}_+^n into \mathbb{R}^n , and agrees with ξ on $X \cap \mathbb{R}_+^n$.

Now extend ξ_β^0 to a C^1 map on all \mathbb{R}^n employing the following recursive procedure; Given ξ^{i-1} defined on $\{\pi \in \mathbb{R}^n \mid \pi_i \geq 0, \dots, \pi_n \geq 0\}$, define ξ^i on $\{\pi \in \mathbb{R}^n \mid \pi_{i+1} \geq 0, \dots, \pi_n \geq 0\}$ by the rule

$$\xi^i(\pi_1, \dots, \pi_n) = \begin{cases} \xi^{i-1}(\pi_1, \dots, \pi_n) & \text{if } \pi_i \geq 0, \\ -\xi^{i-1}(\pi_1, \dots, \pi_{i-1}, -\pi_i, \pi_{i+1}, \dots, \pi_n) \\ + 2\xi^{i-1}(\pi_1, \dots, \pi_{i-1}, 0, \pi_{i+1}, \dots, \pi_n) & \text{if } \pi_i < 0. \end{cases}$$

Note that ξ^i extends ξ^{i-1} and is C^1 . Therefore $\xi^* = \xi^n$ extends ξ_β^0 to all \mathbb{R}^n . Q.E.D.

We shall use this lemma to assume that g is defined on X (although there is no reason to expect that ξ satisfies Assumptions 2 and 3 at points not in $X \cap \mathbb{R}_+^n$).

4. REGULAR ECONOMIES

When studying fixed points of g we want to rule out certain degenerate situations. For this purpose we employ the notion of regular economy introduced by Debreu. References for the technical concepts employed in this and subsequent sections are the books on differential topology by Milnor [25], Guillemin and Pollack [16], and Hirsch [17].

DEFINITION: Consider a C^1 map $f : M \rightarrow N$ from a smooth manifold of dimension m to a smooth manifold of dimension n . A point $x \in M$ is a *regular point* if $Df_x : T(M)_x \rightarrow T(N)_{f(x)}$ has rank n ; in other words, is onto. A point $y \in N$ is a *regular value* if every point x for which $f(x) = y$ is a regular point. Points in M that are not regular points are *critical points*; points in N that are not regular values are *critical values*.

By convention, any point y for which the set $f^{-1}(y)$ is empty is a regular value. Also, if $m < n$, then clearly every point $x \in M$ is a critical point. We extend these concepts to maps such as g that are not everywhere differentiable by requiring that the map Df_x exist at a point x for x to be a regular point.

In the following analysis we focus our attention on economies (ξ, A) for which 0 is a regular value of the map $(g - I): X \rightarrow R^n$ where I is the identity map on R^n . Unfortunately, the map g defined in the previous section is not everywhere differentiable. The projection map p^{S_A} has kinks because of the boundary of S_A . Consequently, g is only piecewise differentiable. As we shall see, however, g is a smooth map on any open set $U \subset X$ such that all points in the image $g(U)$ are contained in the same face of S_A . To ensure that g is differentiable at every equilibrium, let us make the following nondegeneracy assumptions on (ξ, A) :

ASSUMPTION 6: No column of A can be expressed as a linear combination of fewer than n other columns.

ASSUMPTION 7: Let $B(\hat{\pi})$ denote the submatrix of A whose columns are all the activities earning zero profit at π . At every equilibrium $\hat{\pi}$ all \hat{y}_b are strictly positive in the equation $\xi(\hat{\pi}) = \sum_{b \in B(\hat{\pi})} \hat{y}_b b$.

We now turn our attention to finding an expression for $Dg_{\hat{\pi}} - I$ for an equilibrium $\hat{\pi}$ of some economy (ξ, A) satisfying Assumptions 1-7.

LEMMA 2: Let p be the map that projects any point $q \in R^n$ into the nonempty set $\{x | x'C = c'\}$. If the columns of C are linearly independent, then p is a smooth function with Jacobian matrix $Dp_q = I - C(C'C)^{-1}C'$ for all $q \in R^n$.

The proof of this result is a straightforward application of the Kuhn-Tucker theorem. Of course, $I - C(C'C)^{-1}C'$ is simply the orthogonal projection into the null space of C , as anyone familiar with least squares regression techniques would expect.

We want to apply this lemma to $g - I$. Note that, at any fixed point $\hat{\pi}$, $\hat{\pi}'B(\hat{\pi}) = 0$ and Assumption 6 imply that $B(\hat{\pi})$ has fewer than n columns and these are linearly independent. Let $C = [e B(\hat{\pi})]$; the columns of C are linearly independent since $\hat{\pi}'e = 1$ while $\hat{\pi}'B(\hat{\pi}) = 0$.

THEOREM 2: Let (ξ, A) be any economy satisfying Assumptions 1-7. The map g is differentiable in some open neighborhood of every fixed point $\hat{\pi}$, and the Jacobian matrix $Dg_{\hat{\pi}} - I$ equals $(I - C(C'C)^{-1}C')(I + D\xi_{\hat{\pi}}) - I$.

PROOF: By Assumption 7 any activity a^i not in $B(\hat{\pi})$ at equilibrium $\hat{\pi}$ is such that $\hat{\pi}'a^i = g(\hat{\pi})'a^i < 0$. Therefore $\hat{\pi}$ has an open neighborhood U such that $g(\pi)'a^i < 0$ for all $\pi \in U$ by the continuity of g . Lemma 2 implies that, if all the constraints $g(\pi)'B(\hat{\pi}) \leq 0$ are satisfied with equality, then $g(\pi)$ varies smoothly with π . Note, however, that $\hat{y}_b > 0$ for $b \in B(\hat{\pi})$ implies that there is an open

neighborhood $V \subset U$ of $\hat{\pi}$ where $g(\pi)'B(\hat{\pi}) = 0$. Using the chain rule, we can differentiate g to obtain

$$Dg_{\pi} - I = (I - C(C'C)^{-1}C')(I + D\xi_{\pi}) - I$$

for all $\pi \in V$.

Q.E.D.

DEFINITION: An economy $(\xi, A) \in \mathcal{E}$ that satisfies Assumptions 6 and 7 and has $Dg_{\hat{\pi}} - I$ nonsingular at every equilibrium is a *regular economy*. The set of regular economies is denoted $\mathcal{R} \subset \mathcal{E}$. Economies that are not regular are *critical economies*.

To justify this terminology, we note that Theorem 2 implies that 0 is a regular value of $g - I$ if (ξ, A) is a regular economy. As we shall see, the conditions that define a regular economy are satisfied by almost all economies in \mathcal{E} . Of course, the sense of the phrase "almost all" has been made more precise. We shall also demonstrate that these conditions for regularity are equivalent to those given by Debreu [2] for the special case of a pure exchange economy with all equilibria strictly positive.

Another useful concept is that of the equilibrium price correspondence.

DEFINITION: The *equilibrium price correspondence* $\Pi : \mathcal{E} \rightarrow S$ associates with any economy the set of its equilibrium price vectors.

THEOREM 3: Π is an upper-semi-continuous, point-to-set correspondence.

PROOF: Let $(\xi^i, A^i) \rightarrow (\xi, A)$ and $\pi^i \rightarrow \pi$ where $\pi^i \in \Pi(\xi^i, A^i)$. We want to show that $\pi \in \Pi(\xi, A)$. Let us define the production set $Y(A) = \{Ay | y \geq 0\}$. We can rephrase our equilibrium conditions as $\Pi(\xi, A) = \{\pi \in S | \pi'A \leq 0, \xi(\pi) \in Y(A)\}$. Now, if all A^i satisfy Assumptions 4 and 5, $A^i \rightarrow A$ implies $Y(A^i) \rightarrow Y(A)$. (We can use the closed convergence topology for nonempty, closed subsets of R^n to make precise the notion of convergence of sets.) $\xi^i(\pi^i)$ is jointly continuous in ξ^i and π^i , where ξ^i varies in the topology of uniform C^1 convergence on compacta; $\pi^i A^i$ is likewise jointly continuous in π^i and A^i . Therefore we have $\pi^i A^i \rightarrow \pi'A \leq 0$ and $\xi^i(\pi^i) \rightarrow \xi(\pi)$. Since $Y(A^i) \rightarrow Y(A)$ and $\xi^i(\pi^i) \in Y(A^i)$, $\xi(\pi) \in Y(A)$. Thus $\pi \in \Pi(\xi, A)$. *Q.E.D.*

One immediate consequence of this theorem is that for any economy (ξ, A) the set of equilibria $\Pi(\xi, A)$ is a closed subset of the compact set S ; therefore $\Pi(\xi, A)$ is compact. In the case of a regular economy, the inverse function theorem applied to $g - I$ at $\hat{\pi}$ implies that the equilibria are isolated. Thus any economy $(\xi, A) \in \mathcal{R}$ has only a finite number of equilibria.

THEOREM 4: The equilibrium correspondence Π is continuous on \mathcal{R} and the number of equilibria is locally constant.

PROOF: We model our proof after that of Dierker [7, 8] for the pure exchange model. Let (ξ, A) be a regular economy. The set $\Pi(\xi, A) = (g - I)^{-1}(0)$ consists of a finite number of equilibria π^1, \dots, π^k . Each π^i has a neighborhood that is mapped diffeomorphically by $g - I$ onto a neighborhood of 0. We make use of the following result from Dierker [7]:

LEMMA: Let z map a neighborhood of $\pi \in X$ diffeomorphically onto a neighborhood of $0 \in \mathbb{R}^n$ and $z(\pi) = 0$. There exist a neighborhood V of π and $\varepsilon > 0$ such that every C^1 function \tilde{z} maps V diffeomorphically onto some neighborhood of 0 provided the C^1 distance of $\tilde{z}|_V$ and $z|_V$ is less than ε .

It follows that each π^i has a neighborhood V_i such that $(\tilde{g} - I): V_i \rightarrow \mathbb{R}^n$ is a diffeomorphism onto a neighborhood of the origin if \tilde{g} is C^1 close to g on V_i . Let $g = g(\xi, A)$ and $\tilde{g} = g(\tilde{\xi}, \tilde{A})$ for some $(\tilde{\xi}, \tilde{A}) \in \mathcal{E}$. We have no trouble with the differentiability of g since we can choose V_i small enough for $g|_{V_i}$ to be C^1 by Theorem 2. The topology on \mathcal{E} is fine enough for us to find a neighborhood $\mathcal{U} \subset \mathcal{E}$ of (ξ, A) such that $(\tilde{\xi}, \tilde{A}) \in \mathcal{U}$ implies that $\tilde{g}|_{V_i}$ is smooth and had C^1 distance less than ε from $g|_{V_i}$.

Let each V_i be small enough so that $V_i \cap V_{i'} = \emptyset$ for $i \neq i'$. Recall that Π is upper-semi-continuous. The set $\bigcup_{i=1}^k \text{cl}(V_i)$, where $\text{cl}(V_i)$ is the closure of V_i , is a compact subset of X , implying that $\Pi(\tilde{\xi}, \tilde{A}) \subset \bigcup_{i=1}^k V_i$ for all $(\tilde{\xi}, \tilde{A}) \in \mathcal{U}$. Thus the number of equilibria is locally constant. Furthermore, the correspondence Π can be considered as the union of k single-valued functions on \mathcal{U} . They are continuous since Π is upper-semi-continuous. Q.E.D.

When the economy under investigation is regular this theorem provides us with answers to the questions of local uniqueness and continuity posed earlier.

5. THE GLOBAL INDEX THEOREM

We are concerned with fixed points of the map g defined on the manifold X . Although g is not everywhere differentiable, if the economy (ξ, A) satisfies Assumptions 6 and 7 and has locally unique equilibria, we can smooth g without disturbing its fixed points. Although this operation is not strictly necessary to derive the results that follow, carrying it out simplifies our exposition.

LEMMA 3: Let $g: X \rightarrow X$ be defined as previously. If $(\xi, A) \in \mathcal{E}$ satisfies Assumptions 6 and 7 and if the fixed points of g are isolated, then for any $\varepsilon > 0$ there exist a C^1 map $g^*: X \rightarrow X$ and an open set $V \subset X$ containing $\{\pi \in X | g(\pi) = \pi\}$ that satisfy (a) $\|g^*(\pi) - g(\pi)\| \leq \varepsilon$ for all $\pi \in X$; (b) $g^*(\pi) = g(\pi)$ for all $\pi \in V$; (c) $g^*(\hat{\pi}) = \hat{\pi}$ implies $\hat{\pi} \in V$.

PROOF: We make use of the following well known theorem (see, for example, Dieudonne [10, p. 133]):




FIGURE 3

WEIERSTRASS APPROXIMATION THEOREM: *Let $X \subset \mathbb{R}^n$ be a compact set and $g : X \rightarrow \mathbb{R}^n$ be a continuous map. For any $\varepsilon > 0$ there exists a polynomial map $h : X \rightarrow \mathbb{R}^n$ such that $\|h(\pi) - g(\pi)\| \leq \varepsilon$ for all $\pi \in X$.*

The set $\{\pi \in X \mid g(\pi) = \pi\}$ consists of a finite number of points π^1, \dots, π^k . By Theorem 2, for each π^i we can find some $\alpha_i > 0$ such that $g(\pi)$ is differentiable on some open ball $U_i \subset X$ centered at π^i with radius α_i . Choose $\alpha_1, \dots, \alpha_k$ small enough so that $U_i \cap U_{i'} = \emptyset$ for $i \neq i'$. For each π^i choose some β_i such that $\alpha_i > \beta_i > 0$. Let V_i be the open ball centered at π^i with radius β_i . Note that the set $V = \bigcup_{i=1}^k V_i$ contains $\{\pi \in X \mid g(\pi) = \pi\}$ and that g is differentiable on the set $U = \bigcup_{i=1}^k U_i$. Let $\theta^i : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function such that $\theta^i(\pi) = 1$ if $\|\pi^i - \pi\| \leq \beta_i$, $\theta^i(\pi) = 0$ if $\|\pi^i - \pi\| \geq \alpha_i$, and $0 < \theta^i(\pi) < 1$ if $\beta_i < \|\pi^i - \pi\| < \alpha_i$. Let $\theta : \mathbb{R}^n \rightarrow \mathbb{R}$ be the smooth function $\theta(\pi) = \sum_{i=1}^k \theta^i(\pi)$. Note that $\theta(\pi) = 1$ for $\pi \in \text{cl}(V)$, $\theta(\pi) = 0$ for $\pi \in X \setminus U$, and $0 < \theta(\pi) < 1$ for $\pi \in U \setminus \text{cl}(V)$.

We are now ready to construct a map g^* satisfying the conditions of the lemma. The continuous function $\|g(\pi) - \pi\|$ achieves a minimum $\delta > 0$ on the compact set $X \setminus V$. By the above theorem there exists a smooth map $h : X \rightarrow \mathbb{R}^n$ such that $\|h(\pi) - g(\pi)\| \leq \varepsilon < \delta$ for all $\pi \in X$. Since g maps X into S_A , that is, maps X into its interior, we can choose ε small enough so that $h : X \rightarrow X$. Define the map $g^* : X \rightarrow X$ as $g^*(\pi) = \theta(\pi)g(\pi) + (1 - \theta(\pi))h(\pi)$. Note that g^* is C^1 and agrees with g on V . Also note that, for any $\pi \in X \setminus V$, $\|g^*(\pi) - g(\pi)\| \leq \varepsilon$ and $\|g(\hat{\pi}) - \hat{\pi}\| \geq \delta > \varepsilon$ implies that $g(\pi) \neq \pi$. Hence g^* is the map that we desire. Q.E.D.

We now present the formula for computing the local Lefschetz number of an isolated fixed point $\hat{\pi}$ of the map g . Actually, the local Lefschetz number of an isolated fixed point is a purely topological concept, which requires only the continuity of g in order to be defined. However, in keeping with our differentiable approach to the study of equilibria, we shall concern ourselves only with cases where g is differentiable at its fixed points. The above formula indicates that, if $(\xi, A) \in \mathcal{E}$ satisfies Assumptions 6 and 7 and has locally unique equilibria, then, for our purposes, we can assume that $g_{(\xi, A)}$ is C^1 .

For a C^1 map $g : X \rightarrow X$ we compute the local Lefschetz number of an isolated fixed point $\hat{\pi}$ as follows: Take a small closed ball B containing $\hat{\pi}$ and no other fixed point. The rule $v(\pi) = [g(\pi) - \pi] / \|g(\pi) - \pi\|$ defines a smooth map from the boundary of B , ∂B , to the unit sphere, Σ . The local Lefschetz number is defined to be the degree of this map v , which can be any integer. Intuitively, the degree of v measures the number of times v wraps around Σ , taking orientation into account. To be more specific, we choose some regular value y of v and count the number of points in its preimage $v^{-1}(y)$, adding +1 for every point $\pi \in v^{-1}(y)$ such that nonsingular linear map $Dv_\pi : T(\partial B)_\pi \rightarrow T(\Sigma)_y$ preserves orientation and adding -1 for every point $\pi \in v^{-1}(y)$ such that Dv_π reverses orientation. It is well known that this calculation is independent of the regular value y and the ball B and that a regular value always exists (see Hirsch [17, p. 124]).

In the special case where $\hat{\pi}$ is a regular point of $g - I$ this reduces to the following rule (see Hirsch [17, p. 122]): $L_{\hat{\pi}}(g) = +1$ if $Dg_{\hat{\pi}} - I$ preserves orientation. $L_{\hat{\pi}}(g) = -1$ if $Dg_{\hat{\pi}} - I$ reverses orientation. Thus $L_{\hat{\pi}}(g) = \text{sgn}(\det [Dg_{\hat{\pi}} - I])$ when $Dg_{\hat{\pi}} - I$ is nonsingular. Here we define the function $\text{sgn} : \mathbb{R} \rightarrow \{-1, 0, +1\}$ by the rule

$$\text{sgn}(x) = \begin{cases} -1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ +1 & \text{if } x > 0. \end{cases}$$

THEOREM 5: Let $(\xi, A) \in \mathcal{E}$ be any economy that satisfies Assumptions 6 and 7 and has locally unique equilibria. $\sum_{\pi = g(\pi)} L_\pi(g) = (-1)^n$.

PROOF: This theorem is a special case of a major theorem in algebraic topology, the Lefschetz fixed point theorem (see Dold [11] and Guillemin and Pollack [16,

pp. 119–130]). Saigal and Simon [26] prove a version particularly suitable to our purposes. The basic idea is that, if $g : X \rightarrow X$ is a C^1 map with isolated fixed points and if we calculate the local Lefschetz number $L_{\hat{\pi}}(g)$ of a fixed point $\hat{\pi}$ as above, there is a global Lefschetz number $L(g) = \sum_{\pi=g(\pi)} L_{\pi}(g)$ that is a homotopy invariant. In fact, $L(g)$ is independent of g ; it depends only on the manifold X .

Let c be the map $c : X \rightarrow X$ that maps every point into the constant $\bar{x} = (1/n, \dots, 1/n)$. Since X is convex, we are able to construct the homotopy $G : X \times [0, 1] \rightarrow X$ by defining $G(\pi, t) = (1-t)c(\pi) + tg(\pi)$. Note that c is a smooth map and \bar{x} is the unique fixed point of c . Therefore $L(c) = L_{\bar{x}}(c)$. Since $L(g)$ is a homotopy invariant, $L(g) = L(c) = L_{\bar{x}}(c) = \text{sgn}(\det[-I]) = (-1)^n$.

Q.E.D.

Note that this theorem implies the existence of an equilibrium for (ξ, A) . Figure 4 illustrates this result for the case where g maps the unit interval into itself. Here $\sum_{\pi=g(\pi)} L_{\pi}(g) = L(g) = -1$.

Although we can compute the local Lefschetz number of a fixed point whenever it is isolated, we shall have little use for it except when (ξ, A) is a regular economy.




FIGURE 4

DEFINITION: For any economy $(\xi, A) \in \mathcal{E}$ satisfying Assumptions 6 and 7, if an equilibrium $\hat{\pi}$ is such that $[(I - C(C'C)^{-1}C')D\xi_{\hat{\pi}} - I]$ is nonsingular, then *index* $(\hat{\pi})$ is defined as $(-1)^n \text{sgn}(\det [(I - C(C'C)^{-1}C')D\xi_{\hat{\pi}} - I])$.

The advantage of this concept of index is that it can be calculated without reference to the map g . Note that, if (ξ, A) is regular, then $\text{index}(\hat{\pi}) = (-1)^n L_{\hat{\pi}}(g)$.

However, we leave index $(\hat{\pi})$ undefined in cases where $\hat{\pi}$ might not be a regular value of $g - I$.

Let us now state our central result, which is an immediate consequence of Theorem 5 and the definitions of index $(\hat{\pi})$ and regular economy.

THEOREM 6: *If (ξ, A) satisfies Assumptions 1-7 and is a regular economy, then index $(\hat{\pi})$ is defined and equal to +1 or -1 at every equilibrium $\hat{\pi}$ and $\sum_{\pi \in \Pi(\xi, A)} \text{index}(\pi) = +1$.*

6. CALCULATION OF THE INDEX

To apply our results to specific economic models, we need to develop alternative expressions for the index $(\hat{\pi})$. One way to do this is to manipulate the matrix $[D\xi_{\hat{\pi}} - C(C'C)^{-1}C'D\xi_{\hat{\pi}} - C(C'C)^{-1}C']$ without changing the sign of the determinant. Elementary operations with this property can be found in any standard text on linear algebra; our reference is Gantmacher [14]. However, to the writer's knowledge, the following useful result in linear algebra is not found elsewhere.

LEMMA 4: *Let C be an $n \times k$ matrix of full column rank $k \leq n$ and let J be an $n \times n$ matrix. The determinant of $(I - C(C'C)^{-1}C')(I + J) - I$ has the same sign as*

$$\det \begin{bmatrix} J & C \\ C' & 0 \end{bmatrix}.$$

PROOF: $(I - C(C'C)^{-1}C')(I + J) - I = J - C(C'C)^{-1}C'J - C(C'C)^{-1}C'$. The determinant of this matrix equals that of

$$\begin{bmatrix} J - C(C'C)^{-1}C'J - C(C'C)^{-1}C' & 0 \\ C'J + C' & I \end{bmatrix}.$$

Adding the last row pre-multiplied by $C(C'C)^{-1}$ to the first, we do not change the determinant of this matrix. We now have

$$\det \begin{bmatrix} J & C(C'C)^{-1} \\ C'J + C' & I \end{bmatrix}.$$

Subtracting the first row pre-multiplied by C' from the last, we are left with

$$\det \begin{bmatrix} J & C(C'C)^{-1} \\ C' & 0 \end{bmatrix}.$$

Note that $C'C$ is positive definite and hence has a positive determinant. The sign of the above expression therefore remains unchanged when the matrix is post-multiplied by

$$\begin{bmatrix} I & 0 \\ 0 & C'C \end{bmatrix}$$

where I is $n \times n$:

$$\det \left(\begin{bmatrix} J & C(C'C)^{-1} \\ C' & 0 \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & C'C \end{bmatrix} \right) = \det \begin{bmatrix} J & C \\ C' & 0 \end{bmatrix}. \quad Q.E.D.$$

We can use this lemma to calculate index ($\hat{\pi}$) of any regular equilibrium $\hat{\pi}$ of an economy that satisfies Assumptions 1-7:

$$\begin{aligned} \text{index}(\hat{\pi}) &= (-1)^n \text{sgn} \left(\det \begin{bmatrix} D\xi_{\hat{\pi}} & e' & B(\hat{\pi}) \\ e & 0 & 0 \\ B'(\hat{\pi}) & 0 & 0 \end{bmatrix} \right) \\ &= (-1)^n \text{sgn} \left(\det \begin{bmatrix} 0 & e' & 0 \\ e & D\xi_{\hat{\pi}} & B(\hat{\pi}) \\ 0 & B'(\hat{\pi}) & 0 \end{bmatrix} \right). \end{aligned}$$

An alternative expression is

$$\text{index}(\hat{\pi}) = \text{sgn} \left(\det \begin{bmatrix} -D\xi_{\hat{\pi}} + E & -B(\hat{\pi}) \\ B'(\hat{\pi}) & 0 \end{bmatrix} \right)$$

where E is an $n \times n$ matrix whose every element is unity. To see why this is so, note that

$$\det \begin{bmatrix} -D\xi_{\hat{\pi}} + E & -B(\hat{\pi}) \\ B'(\hat{\pi}) & 0 \end{bmatrix} = (-1)^n \det \begin{bmatrix} 1 & e' & 0 \\ 0 & D\xi_{\hat{\pi}} - E & B(\hat{\pi}) \\ 0 & B'(\hat{\pi}) & 0 \end{bmatrix}.$$

If we post-multiply the second column of the final matrix by $\hat{\pi}$ and subtract from the first column, we do not change the determinant. The homogeneity assumption (Assumption 2), when differentiated, implies that $D\xi_{\hat{\pi}}\pi = 0$ for any $\pi \in R_+^n \setminus \{0\}$. Also $e'\hat{\pi} = 1$ and $B'(\hat{\pi})\hat{\pi} = 0$. Thus we are left with

$$(-1)^n \det \begin{bmatrix} 0 & e' & 0 \\ e & D\xi_{\hat{\pi}} - E & B(\hat{\pi}) \\ 0 & B'(\hat{\pi}) & 0 \end{bmatrix} = (-1)^n \det \begin{bmatrix} 0 & e' & 0 \\ e & D\xi_{\hat{\pi}} & B(\hat{\pi}) \\ 0 & B'(\hat{\pi}) & 0 \end{bmatrix}.$$

Let the matrix \bar{J} be formed by deleting from $D\xi_{\hat{\pi}}$ all rows and columns i for which $\hat{\pi}_i = 0$ and then deleting any one more row and column. Let \bar{B} be formed by deleting from $B(\hat{\pi})$ the columns corresponding to disposal activities, all rows i for which $\hat{\pi}_i = 0$, and the same additional row as that deleted from $D\xi_{\hat{\pi}}$. Note that Walras's law (Assumption 3), when differentiated, implies that $(D\xi_{\hat{\pi}})'\hat{\pi} = -\xi(\hat{\pi}) = -B(\hat{\pi})y_B$. Using this observation and matrix manipulations similar to those above, we can easily demonstrate that

$$\text{index}(\hat{\pi}) = \text{sgn} \left(\det \begin{bmatrix} -\bar{J} & -\bar{B} \\ \bar{B}' & 0 \end{bmatrix} \right)$$

if more than one $\hat{\pi}_i$ is strictly positive; $\text{index}(\hat{\pi}) = +1$ otherwise.

Now, if $A = -I$ (the case of the pure exchange model) and if $\xi_i(\pi) > 0$ for any π such that $\pi_i = 0$, then any equilibrium $\hat{\pi}$ is strictly positive and $\text{index}(\hat{\pi}) = \text{sgn}(\det[-\bar{J}])$. Such is the situation analyzed by Dierker [6, 7, 8] and Varian [32], who calculate the same expression for the index. Moreover, since Assumptions 6 and 7 are satisfied by any economy $(\xi, -I)$, if $\det[-\bar{J}] \neq 0$ at any equilibrium, then $(\xi, -I)$ is regular. This concept of regularity is therefore equivalent to that of Debreu [2]. Thus this formulation of $\text{index}(\hat{\pi})$ allows us to compare our results with those previously known for pure exchange economies. Even for these economies our results are more general than those given elsewhere; we allow some prices to be zero at equilibrium.

One other point is worth mentioning. For the sake of clarity, we have defined the map g on the n manifold X . However, since $g(X) \subset S_A$, we could have well worked with the $n-1$ manifold $X \cap \{\pi \in \mathbb{R}^n | \sum_{i=1}^n \pi_i = 1\}$. Many writers find it more elegant to work with an $n-1$ manifold in this type of model, identifying S with its natural projection into the last $n-1$ price coordinates. Although this is less trivial than it might seem, we can use the homogeneity assumption (Assumption 2) and Walras's law (Assumption 3) to drop the first coordinate from consideration. In this setting the natural definition for $\text{index}(\hat{\pi})$, at least when $\hat{\pi}$ is strictly positive, becomes $(-1)^{n-1} \text{sgn}(\det[\bar{J} - \bar{B}(\bar{B}'\bar{B})^{-1}\bar{B}'\bar{J} - \bar{B}(\bar{B}'\bar{B})^{-1}\bar{B}'])$. Rather than going through the derivation of this, which requires some work, we simply note that, by Lemma 4, this is equivalent to the expression for the index that we have already derived.

7. GENERICITY OF REGULAR ECONOMIES

The value of the global index theorem depends on how common a situation regularity is in the space of economies. Intuitively, we view economies that are not regular, critical economies, as somehow degenerate. Thus we might hope that such economies form a very small subset of \mathcal{E} . In fact this is the case.

DEFINITION: A property that holds for some subset \mathcal{V} of a topological space \mathcal{U} is a *generic property* of \mathcal{U} if \mathcal{V} is open and dense in \mathcal{U} .

We shall argue that the property of regularity is generic, in other words, that \mathcal{R} is open and dense in \mathcal{E} .

It should be noted that genericity as defined above is stronger than the usual concept. Mathematicians often speak of some subset \mathcal{V} of a space \mathcal{U} as being residual if it contains the intersection of a countable number of open dense sets. For example, the set of irrational numbers is a residual subset of the reals. A property that holds for a residual subset of \mathcal{U} is then called a generic property. An open dense set is, of course, residual, but a residual set, although dense, is not necessarily open. We shall insist on the stronger definition, however. This is to ensure that the set of critical economies is not dense in \mathcal{E} . The set of irrational numbers is not open; its complement, the set of rational numbers, is dense on the real line.

The difficult half of proving that regularity is a generic property of \mathcal{E} is proving that \mathcal{R} is dense in \mathcal{E} ; Theorem 4 already implies that it is open. Our approach can be motivated by an argument reminiscent of the counting of equations and variables by Walras. Our equilibrium conditions can be expressed as a system of $n + k$ variables in $n + k + 1$ equations

$$\begin{aligned} B' \hat{\pi} &= 0, \\ \xi(\hat{\pi}) &= B\hat{y}, \\ e' \hat{\pi} &= 1, \end{aligned}$$

where B is the $n \times k$ matrix of activities $B(\hat{\pi})$. Walras's law implies that one of the n equations in $\xi(\hat{\pi}) = B\hat{y}$ can be eliminated. The conditions that ensure regularity also ensure that the equilibrium pair $(\hat{\pi}, \hat{y})$ is a locally unique solution to this system that is stable under small perturbations. If either Assumption 6 or the non-zero determinant condition does not hold, we may not have enough independent equations for the equilibrium to be totally unique. If, on the other hand, Assumption 7 does not hold, then we may have too many independent equations for a solution to exist. If we do not have as many independent equations as variables, then it is intuitively plausible that some very slight perturbation in the underlying parameters of the economy could make the equations independent. Similarly, if there are not as many independent variables as equations, then some slight perturbation of the system could make a solution impossible. What we need is freedom to make perturbations in a sufficient number of directions.

The proof that the set of matrices satisfying Assumption 6 is open and dense in \mathcal{A} is trivial; we omit it here. We can demonstrate that \mathcal{R} is dense in \mathcal{E} by proving that for any economy (ξ^0, A) that satisfies Assumption 6, but not Assumption 7 or not the non-zero determinant condition, there is another economy (ξ^1, A) that is regular with $d_S(\xi^0, \xi^1) < \varepsilon$ for any $\varepsilon > 0$. We reduce the problem from one in the infinite dimensional vector space \mathcal{D} to the finite dimensional vector space R^n by parameterizing any excess demand function $\xi \in \mathcal{D}$ with a perturbation vector $v \in R^n$. Define the function $\delta(\pi, v) = (\delta_1(\pi, v), \dots, \delta_n(\pi, v))$,

$$\delta_i(\pi, v) = \frac{\sum_{l=1}^n \pi_l v_l}{\sum_{l=1}^n \pi_l} - v_i \quad (i = 1, \dots, n).$$

δ satisfies Assumptions 1–3, and therefore so does $\xi + \delta$ if ξ does. For any economy (ξ, A) we consider the family of economies $\{(\xi_v, A) \in \mathcal{E} \mid \xi_v(\pi) = \xi(\pi) + \delta(\pi, v), v \in R^n\}$. Our goal is to demonstrate that the set of regular economies is dense in this n parameter family. If it is, the topology on \mathcal{E} is such that for any $\varepsilon > 0$ we can find a $\varepsilon' > 0$ such that $\|v\| < \varepsilon'$ implies that $d[(\xi, A), (\xi_v, A)] < \varepsilon$. Consequently, if regular economies are dense in this n parameter family, they are dense in \mathcal{E} .

In R^n the concept of density is related to that of Lebesgue measure. Recall that a set of $U \subset R^n$ has Lebesgue measure zero if it is possible to cover U with a countable number of rectangular solids with arbitrarily small volume. The complement in R^n of a set with Lebesgue measure zero has full Lebesgue measure. This concept of Lebesgue measure is easily extended to manifolds using local parameterizations (see Guillemin and Pollack [16, pp. 204–205]). It is well known that a set of full Lebesgue measure is dense (see Milnor [26, pp. 10–11]). A theorem in differential topology that is a direct consequence of Sard's theorem is particularly suited to our purpose (Guillemin and Pollack [16, pp. 67–69]).

TRANSVERSALITY DENSITY THEOREM: *Let $M, V,$ and N be smooth manifolds without boundary where $\dim M = m, \dim N = n,$ and $m \leq n,$ and let $y \in N.$ Suppose that $F: M \times V \rightarrow N$ is a C^1 map such that, for every $(x, v) \in M \times V,$ $\text{rank } DF_v(x, v) = n.$ The set of $v \in V$ for which $F(x, v) = y$ implies that $\text{rank } DF_x(x, v) = n$ has full Lebesgue measure.*

Before applying this theorem to our problem, let us develop a few preliminary concepts. For any $n \times k$ matrix $B, 0 \leq k \leq n - 1,$ let

$$K_B = \{x \in R^n \mid B'x = 0, e'x = 1\},$$

$$O_B = \{x \in R^n \mid B'x = 0, e'x = 0\}.$$




FIGURE 5

Note that, at any equilibrium $\hat{\pi}$ of an economy $(\xi, A) \in \mathcal{E},$ both $K_{B(\hat{\pi})}$ and $K_{B(\hat{\pi})} \cap \text{int } X$ are smooth and without boundary. Consider the C^1 map $f^{B(\hat{\pi})}: (K_{B(\hat{\pi})} \cap \text{int } X) \rightarrow O_{B(\hat{\pi})}$ defined by the rule $f^{B(\hat{\pi})}(\pi) = p^{K_{B(\hat{\pi})}}(\pi + \xi(\pi)) - \pi = p^{O_{B(\hat{\pi})}}(\xi(\pi)).$ If (ξ, A) satisfies Assumptions 6 and 7, then $f^{B(\hat{\pi})}$ agrees with $g - I$ in

some neighborhood of $\hat{\pi}$ on $K_{B(\hat{\pi})}$. At an equilibrium $\hat{\pi}$, $f^{B(\hat{\pi})}(\hat{\pi}) = 0$. The derivative map $Df_{\hat{\pi}}^{B(\hat{\pi})} : T(K_{B(\hat{\pi})} \cap \text{int } X)_{\hat{\pi}} \rightarrow T(O_{B(\hat{\pi})})_0$ maps $O_{B(\hat{\pi})}$ into itself; on $O_{B(\hat{\pi})}$, $Df_{\hat{\pi}}^{B(\hat{\pi})} = (I - C(C'C)^{-1}C')D\xi_{\hat{\pi}}$.

For any economy (ξ, A) we focus our attention on all possible maps f^B where B is some $n \times k$ submatrix of A , $0 \leq k \leq n - 1$, such that $K_B \cap S_B \neq \emptyset$. Obviously, every equilibrium $\hat{\pi}$ is such that $\hat{\pi} \in K_B$ and $f^B(\hat{\pi}) = 0$ for some such B , namely $B(\hat{\pi})$. The converse, however, does not necessarily hold; in addition to $\hat{\pi} \in K_B$ and $f^B(\hat{\pi}) = 0$ the condition $\hat{\pi} \in S_A$ is needed for $\hat{\pi}$ to be an equilibrium. We now parameterize our maps f^B with the perturbation vector $v \in R^n$ by defining the C^1 map $F^B : (K_B \cap \text{int } X) \times R^n \rightarrow O_B$ by the rule $F^B(\pi, v) = \rho^{O_B}(\xi(\pi) + \delta(\pi, v))$.

To apply the transversality density theorem, we must demonstrate that for all $(\pi, v) \in (K_B \cap \text{int } X) \times R^n$ the derivative map $DF_v^B : R^n \rightarrow O_B$ has rank $n - k - 1$, in other words, is onto. $DF_v^B(\pi, v) = (I - C(C'C)^{-1}C')D\delta_v$ where $C = [e \ B]$. Now

$$D\delta_v(\pi, v) = \begin{bmatrix} \pi_1 - 1 & \dots & \pi_n \\ \pi_1 & \dots & \pi_n - 1 \end{bmatrix}$$

has rank $n - 1$ since $D\delta_v(\pi, v)e = 0$ but the $(n - 1) \times (n - 1)$ matrix formed by deleting any row and column j for which $\pi_j > 0$ is nonsingular. Letting $\rho = \text{rank}(I - C(C'C)^{-1}C')D\delta_v$, we note that

$$\begin{aligned} \rho + k + 1 &= \text{rank} \begin{bmatrix} (I - C(C'C)^{-1}C')D\delta_v & C \\ 0 & I \end{bmatrix} \\ &= \text{rank} \begin{bmatrix} D\delta_v & C \\ (C'C)^{-1}C'D\delta_v & I \end{bmatrix} \\ &= \text{rank} \begin{bmatrix} D\delta_v & C \\ 0 & 0 \end{bmatrix} \\ &= \text{rank}[D\delta_v \ e \ B]. \end{aligned}$$

Clearly, $\text{rank}[D\delta_v \ e \ B] \geq n$ since $D\delta_v$ has rank $n - 1$, $D\delta_v e = 0$, and $e'e = n$. However, $(I - C(C'C)^{-1}C')D\delta_v$ maps into O_B , implying that $\rho = n - k - 1$.

We first demonstrate that, for almost all $v \in R^n$, the economy (ξ_v, A) satisfies Assumption 7. Let B^* be any $n \times k^*$, $n \geq k^* \geq k$, submatrix of A that has B as a submatrix. Obviously, there are only a finite number of such combinations of matrices B and B^* for any economy (ξ, A) . $K_{B^*} \cap \text{int } X$ is an $n - k^* - 1$ submanifold of the $n - k - 1$ manifold $K_B \cap \text{int } X$. We restrict F^B to domain $(K_{B^*} \cap \text{int } X) \times R^n$. As a consequence of the transversality density theorem, for almost all $v \in R^n$, $F^B(\pi, v) = 0$ implies that $\text{rank } DF_{\pi}^B = n - k - 1$. However, $k^* > k$ implies $\text{rank } DF_{\pi}^B \leq n - k^* - 1 < n - k - 1$. Thus, for almost all $v \in R^n$, there is no $\pi \in K_{B^*} \cap \text{int } X$ such that $F^B(\pi, v) = 0$. Intuitively, although F^B takes $(K_{B^*} \cap \text{int } X) \times R^n$ into O_B , the image $F^B(K_{B^*} \cap \text{int } X, v)$ is a very small subset of O_B for any fixed $v \in R^n$. Indeed, for a set U of full Lebesgue measure in R^n , the image $F^B(K_{B^*} \cap \text{int } X, U)$ does not contain 0. We can repeat this same argument for all possible

combinations B and B^* . The intersection of a finite number of sets of full Lebesgue measure also has full Lebesgue measure.

Now, if we apply the transversality density theorem to $F^B : (K_B \cap \text{int } X) \times R^n \rightarrow O_B$, we establish that, for almost all $v \in R^n$, $F^B(\pi, v) = 0$ implies that $\text{rank } DF_{\pi}^B = n - k - 1$. Thus, for almost all $v \in R^n$, if $\hat{\pi}$ is an equilibrium of (ξ_v, A) , then $DF_{\hat{\pi}}^{B(\hat{\pi})} = (I - C(C'C)^{-1}C')D\xi_{\hat{\pi}}x$ has rank $n - k - 1$. Let $x \in R^n$ be such that

$$(I - C(C'C)^{-1}C')D\xi_{\hat{\pi}}x - C(C'C)^{-1}C'x = 0.$$

Since the columns of $(I - C(C'C)^{-1}C')D\xi_{\hat{\pi}}$ and those of $C(C'C)^{-1}C'$ are orthogonal and $\text{rank } C(C'C)^{-1}C' = k + 1$,

$$\{x \in R^n | (I - C(C'C)^{-1}C')D\xi_{\hat{\pi}}x = 0\} \cap \{x \in R^n | C(C'C)^{-1}C'x = 0\} = \{0\}.$$

Pre-multiplying the above equation by $(I - C(C'C)^{-1}C')$ gives us

$$(I - C(C'C)^{-1}C')D\xi_{\hat{\pi}}x = 0.$$

On the other hand, pre-multiplying by $C(C'C)^{-1}C'$ produces

$$C(C'C)^{-1}C'x = 0.$$

Together these imply that $[(I - C(C'C)^{-1}C')D\xi_{\hat{\pi}} - C(C'C)^{-1}C']$ is nonsingular. Again using the fact that the intersection of a finite number of sets of full Lebesgue measure also has full Lebesgue measure, we have proven the following theorem.

THEOREM 6: *The set of regular economies \mathcal{R} is open and dense in \mathcal{E} .*

8. EXTENSIONS AND CONCLUSIONS

The above treatment of genericity of regular economies is rough in the sense that the perturbation of excess demand functions, while entirely natural for economies where all commodities enter into consumers' final demands, is not always appropriate in economies where production plays an important role. In such economies there are likely to be primary commodities which are inelastically supplied as inputs to the production process and intermediate commodities which are only produced in order to produce other commodities. It is possible to extend our argument to such situations, but there are several minor technical problems. First, we must explicitly deal with the possibility of excess demand being unbounded at some points on the boundary of R_+^n in order to deal properly with primary commodities. Second, we must slightly alter our definition of regularity to deal with the possibility of prices of intermediate commodities being undefined at equilibria where no production takes place.

Another direction in which our discussion can be extended is production technologies with smooth production functions. It is relatively easy to exploit the properties of the profit functions in such a technology to show that our mapping and index theorem carry over when we substitute for activities the input-output vectors of the production functions used at equilibrium.

These matters are discussed in more detail in the writer's doctoral dissertation (Kehoe [18]). Also discussed there are conditions that may be placed on the structure of an economic model such that the index theorem implies uniqueness of equilibrium. We remark briefly on two sets of such conditions, one dealing with the production side of the economy alone, the other dealing with the consumption side alone. First, if we can impose conditions on (ξ, A) that ensure that there are always $n - 1$ activities in use at equilibrium, then

$$\text{index}(\hat{\pi}) = \text{sgn} \left(\det \begin{bmatrix} -\bar{J} & -\bar{B} \\ \bar{B}' & 0 \end{bmatrix} \right) = \text{sgn}(\det[\bar{B}'\bar{B}]) = +1$$

at every equilibrium. The global index theorem then implies that there is a unique equilibrium. Recall that the conditions of the nonsubstitution theorem of input-output analysis imply that there are always $n - 1$ activities in use at equilibrium. Second, if the excess demand function ξ is such that \bar{J} is always negative definite, then there is a unique equilibrium if the production technology is such that (ξ, A) satisfies Assumptions 4-7. This can be seen by noting that

$$\text{index}(\hat{\pi}) = \text{sgn} \left(\det \begin{bmatrix} -\bar{J} & -\bar{B} \\ \bar{B}' & 0 \end{bmatrix} \right) = \text{sgn}(\det[-\bar{J}] \det[-\bar{B}'\bar{J}^{-1}\bar{B}]) = +1$$

at every equilibrium. It is interesting to note that it has been shown by Kihlstrom, Mas-Colell, and Sonnenschein [19] that negative definiteness of \bar{J} implies that the weak axiom of revealed preference holds for ξ , a condition previously known to imply uniqueness.

A final remark should also be made with regard to the necessity of the conditions we have established for uniqueness of equilibrium. Conditions such as gross substitutability in ξ are neither necessary nor sufficient for $\text{index}(\hat{\pi})$ to be positive. Constructing an economy with gross substitutes and with an equilibrium with $\text{index}(\hat{\pi}) = -1$ is fairly easy. Such an equilibrium cannot be unique. This observation suggests that nonuniqueness of equilibrium may be a less pathological situation than is sometimes thought.

Wesleyan University

Manuscript received February, 1979; revision received September, 1979.

REFERENCES

- [1] ARROW, K., AND F. HAHN: *General Competitive Analysis*. San Francisco: Holden-Day, Inc., 1971.
- [2] DEBREU, G.: "Economies with a Finite Set of Equilibria," *Econometrica*, 38 (1970), 387-393.
- [3] ———: "Smooth Preferences," *Econometrica*, 40 (1972), 603-612.
- [4] ———: "Corrigendum," *Econometrica*, 43 (1975), 831-832.
- [5] ———: "Regular Differentiable Economies," *American Economic Review, Papers and Proceedings*, 66 (1976), 280-287.
- [6] DIERKER, E.: "Two Remarks on the Number of Equilibria of an Economy," *Econometrica*, 40 (1972), 951-953.

- [7] ———: *Topological Methods in Walrasian Economics*, Lecture Notes in Economics and Mathematical Systems, 92. New York: Springer-Verlag, 1974.
- [8] ———: "Regular Economies," in *Handbook of Mathematical Economics*, ed. by K. Arrow and M. Intriligator. Amsterdam: North-Holland, forthcoming.
- [9] DIERKER, E., AND H. DIERKER: "The Local Uniqueness of Equilibria," *Econometrica*, 40 (1972), 867-881.
- [10] DIEUDONNE, J.: *Foundations of Modern Analysis*. New York: Academic Press, 1960.
- [11] DOLD, A.: "Fixed Point Index and Fixed Point Theorem for Euclidean Neighborhood Retracts," *Topology*, 4 (1965), 1-8.
- [12] EAVES, B., AND H. SCARF: "The Solution of Systems of Piecewise Linear Equations," *Mathematics of Operations Research*, 1 (1976), 1-27.
- [13] FUCHS, G.: "Private Ownership Economics with a Finite Number of Equilibria," *Journal of Mathematical Economics*, 1 (1974), 141-158.
- [14] GANTMACHER, F.: *The Theory of Matrices*, Volume I, translated from the Russian by K. Hirsch. New York: Chelsea Publishing Company, 1959.
- [15] GREENBERG, M.: *Lectures on Algebraic Topology*. New York: W. A. Benjamin, Inc., 1967.
- [16] GUILLEMIN, Y., AND A. POLLACK: *Differential Topology*. Englewood Cliffs, New Jersey: Prentice Hall, Inc., 1974.
- [17] HIRSCH, M.: *Differential Topology*. New York: Springer-Verlag, 1976.
- [18] KEHOE, T.: "Regularity and Index Theory for Economic Equilibrium Models," Ph.D. Dissertation, Yale University, 1979.
- [19] KIHNSTROM, R., A. MAS-COLELL, AND H. SONNENSCHNEIN: "The Demand Theory of the Weak Axiom of Revealed Preference," *Econometrica*, 44 (1976), 971-978.
- [20] MCFADDEN, D., A. MAS-COLELL, R. MANTEL, AND M. RICHTER: "A Characterization of Community Excess Demand Functions," *Journal of Economic Theory*, 9 (1974), 361-374.
- [21] MAS-COLELL, A.: "Continuous and Smooth Consumers: Approximation Theorems," *Journal of Economic Theory*, 8 (1974), 305-336.
- [22] ———: "On the Continuity of Equilibrium Prices in Constant-Return Production Economies," *Journal of Mathematical Economics*, 2 (1975), 21-33.
- [23] ———: "The Theory of Economic Equilibria from the Differentiable Point of View," Course Notes, Department of Mathematics, University of California, Berkeley, 1976.
- [24] ———: "An Introduction to the Differentiable Approach to the Theory of Economic Equilibria," in *Studies in Mathematical Economics*, ed. by S. Reiter. Washington, D.C.: Mathematical Association of America, forthcoming.
- [25] MILNOR, J.: *Topology from the Differentiable Viewpoint*. Charlottesville, Virginia: University Press of Virginia, 1976.
- [26] SAIGAL, R., AND C. SIMON: "Generic Properties of the Complementarity Problem," *Mathematical Programming*, 4 (1973), 324-335.
- [27] SMALE, S.: "Global Analysis and Economics, IV; Finiteness and Stability of Equilibria with General Consumption Sets and Production," *Journal of Mathematical Economics*, 1 (1974), 119-127.
- [28] ———: "Dynamics in General Equilibrium Theory," *American Economic Review, Papers and Proceedings*, 66 (1976), 288-294.
- [29] ———: "Global Analysis and Economics," in *Handbook of Mathematical Economics*, ed. by K. Arrow and M. Intriligator. Amsterdam: North-Holland, forthcoming.
- [30] SPIVAK, M.: *Calculus on Manifolds*. Menlo Park, California: The Benjamin-Cummings Publishing Company, 1965.
- [31] TODD, M.: "A Note on Computing Equilibria in Economies with Activity Analysis Model of Production," CORE Discussion Paper 7736, Université Catholique de Louvain, 1977.
- [32] VARIAN, H.: "A Third Remark on the Number of Equilibria of an Economy," *Econometrica*, 43 (1974), 985-986.